

AD-A135 851

THE CHEMICAL CHARACTERISTICS OF PGBX(U) MASSACHUSETTS
INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY K BIEMANN
18 AUG 83 N00014-78-C-0421


1/1

UNCLASSIFIED

F/G 6/15

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A135851

12

Final Progress Report

Project Title:

THE CHEMICAL CHARACTERISTICS OF PGB_x

Principal Investigator:

Klaus Biemann, Ph.D.
Professor of Chemistry

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, MA 02139

Contract N00014-78-C-0421

Date of Report:
August 18, 1983

DTIC
ELECTE
DEC 15 1983
S D

DTIC FILE COPY

DISTRIBUTION STATEMENT A

Approved for public release;
Distribution Unlimited

83 09 02 002

Final Progress Report

Title of Project: THE CHEMICAL CHARACTERISTICS OF PGB_x

Principal Investigator: Klaus Biemann, Dept. of Chemistry
Massachusetts Institute of Technology
Cambridge, MA 02139

Contract N00014-78-C-0421

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By <u>Per Ltr. on File</u>	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
<u>A/1</u>	

Objectives


It was first attempted to separate PGB_x , a biologically active material obtained upon alkali treatment of 15-keto-prostaglandin- B_1 (1), into its components, determine their molecular weight and, finally, molecular structure. When it became clear that PGB_x is too complex a mixture, model studies were begun.

To investigate the mechanism by which the chemically active moiety in 15-keto-prostaglandin- B_1 oligomerizes to PGB_x under alkaline conditions, the condensation of 3-ethyl-4-(3'-ketopenten-1-yl) cyclopent-2-enone (2) was investigated.


Summary of Results

A high pressure liquid chromatograph was used to separate various batches of PGB_x and its precursors. Partial separation has been achieved. Field desorption mass spectrometry of PGB_x was attempted but without success. ^{252}Cf plasma desorption mass spectrometry seems to indicate the presence of a mixture ranging from tetramers to octamers of 15-keto- PGB_1 .

The condensation of modified models of the diene-dione moiety (2) of 15-keto- PGB_1 (1) was then studied. It has been found that a number of dimers are involved in this reaction. Furthermore, the results indicate that the anion at C-16 is very reactive and probably the initial step. Gas chromatography (GC), high performance liquid chromatography (HPLC), field desorption (FD), fast atom bombardment (FAB) mass spectrometry, ^1H and ^{13}C -NMR were used for the analysis of the resulting mixtures of isomers and oligomers.



(1)




(2)


Conclusions

The base catalyzed condensation of the model ketone (2), when car-

ried out under conditions somewhat milder than those which convert $\text{PGB}_1(1)$ to PGB_x , produced a number of dimeric compounds. Upon separation by gas and liquid chromatography their structures (3-6) were determined by spectroscopic means and found to be most likely represented by the following:



The presence of isomer (7) is not excluded.



The fact that the simpler model, under mild conditions, can undergo condensation in four or five ways indicates that the oligomer (PGB_x) obtained when treating PGB_1 with alkali must be a very complex mixture. Under those circumstances it is not surprising that the oligomer could not be separated into the individual components.

Just when this work was completed a paper appeared by G.L. Nelson and G.L. Verdine (Tetrahedron Letters, 23, v.1967-1970 (1982)) describing essentially the same results. Although the senior author of that paper (G.L.N.) had agreed to keep the Principal Investigator (K.B.) informed about his work to avoid duplication and to publish the results jointly or simultaneously, this was not done. We therefore decided not to publish our results separately later. This episode is typical of the lack of cooperation and communication that plagued the entire PGB_x project from the outset.

Technical Reports; None

Publications; Optimization of Experimental Procedures for Fast Atom Bombardment Mass Spectrometry. S.A. Martin, C.E. Costello, and K. Biemann, Anal. Chem. 54, 2362-2368 (1982).

8/18/83

LMED
-8